

## MEMORANDUM

To: USACE Colonel James L. Booth, Major Cory Bell, Richard McMillen, SFWMD Governing Board,  
Executive Director Drew Bartlett, Jennifer Reynolds, Lawrence Glenn, DEP Secretary Shawn Hamilton

From: Periodic Scientists Conference Call Participants  
Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex  
Holly Milbrandt & Dana Dettmar - City of Sanibel  
Harry Phillips & Maya Robert - City of Cape Coral  
Allie Pecenka, Rick Bartleson PhD & Matt Depaolis- Sanibel-Captiva Conservation Foundation  
In coordination with Lee County

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **June 4- 10, 2024**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

**Caloosahatchee Conditions Summary:** Flow to the Caloosahatchee Estuary had a 7-day average of **1,690 cfs** at **S-79** with a 7-day average of **1,575 cfs (93%)** coming from the lake at **S-77**. **The 14-day moving average flow at S-79 is 1,998 cfs and has been in the optimum flow envelope** (750 – 2,100 cfs) for **56 days**. The 14-day average flow at S-77 was **1,857 cfs**.

**Recommendation:** On April 13, The Army Corps began targeting a 7-day average pulse release schedule of 2,000 cfs at W.P. Franklin Lock and Dam (S-79) and 0 cfs St. Lucie Lock and Dam (S-80). With the onset of the rainy season and predictions for increased Atlantic storm intensity in the upcoming hurricane season, we ask the Army Corps to remain reactive to changing conditions in Lake Okeechobee and the Caloosahatchee River and estuary to support the ecological health of this system. In addition, we request a pause in releases from Lake Okeechobee to the CRE for as long as basin runoff from prolonged rainfall in Southwest Florida pushes these flows above RECOVER targets.

**USACE Action:** With Lake Okeechobee stage within the Low Sub-band, the Tributary Hydrologic Conditions in the Dry category, Part D of the 2008 LORS suggests "S-79 up to 450 cfs and S-80 up to 200 cfs." On 4/13/24 the USACE began targeting a 7-day average pulse release schedule of 2,000 cfs at W.P. Franklin Lock and Dam (S-79) and 0 cfs St. Lucie Lock and Dam (S-80).

**Lake Flows:** In the past 7 days the total outflow from Lake Okeechobee was **44,504 AF\*** with **21,843 AF** to the Caloosahatchee through **S-77**, **321 AF** to the St. Lucie canal through **S-308**, **939 AF** through the **L8 canal**, and **21,401 AF** to the EAA through **S-351**, **S-352**, and **S-354\***. The total net inflow to the Lake was **3,076 AF (3,076 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1). Water conservation areas received flows of **0 AF**, **1,617 AF**, and **759 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **69 AF**.  
\*Data missing from S-78 on 6/10, S-310 from 6/4- 6/10, S-354 from 6/9- 6/10 and S-80 from 6/7- 6/8 and on 6/10.

**Lake Level: 12.63 ft (Low Sub-Band)**

**Last Week: 12.86 ft**

**Last Year: 14.04 ft**

**7-Day Lake Recession Rate: -0.23 ft/week**

**Lake Okeechobee Inflow: 206 cfs**

**Lake Okeechobee Outflow: 622 cfs**


**Weekly Rainfall Total: WP Franklin: 0.24"**

**Ortona: 4.39"**

**Moore Haven: 2.99"**

**Cyanobacteria Status:** On 6/10/24, sampling for cyanobacteria by the Lee County Environmental Lab reported **moderately abundant** *Dolichospermum*, *Microcystis* and cyano filaments at the **Alva Boat Ramp** as streaks. *Dolichospermum*, *Microcystis*, *Aphanizomenon*, and cyano filaments were **moderately abundant** upstream of the **Franklin Locks** as streaks with heavy accumulation along the Lock and **present** at the **Davis Boat Ramp** as streaks.

**Red Tide:** On 6/7/24, the FWC reported that the red tide organism, *Karenia brevis*, was not observed in samples collected statewide over the past week.



Site	Light Penetration		Turbidity	Target Values
	25% I <sub>z</sub>	Target Values		
	meters		NTU	
Fort Myers	0.9	> 1	4.3	< 18
Shell Point	1.3	>2.2	1.1	< 18
Causeway	4.4	> 2.2	2.2	< 5

25% I<sub>z</sub> is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

**Upper Estuary Conditions:** The 30-day average surface salinity at the Fort Myers Yacht Basin was 3.6 psu, within the suitable range for tape grass.

**Lower Estuary Conditions:** The average salinity at Shell Point RECON was 27 psu, in the optimal range for oysters and seagrass.

**Water Quality Conditions:**

Monitor Site	Salinity (psu) <sup>a</sup> [previous week]	Diss O <sub>2</sub> (mg/L) <sup>b</sup>	FDOM (qsde) <sup>c</sup>	Chlorophyll (µg/L) <sup>d</sup>	Temperature (°F)
Beautiful Island	0.2 – 0.5 [0.2 – 0.2]	1.3 – 7.5	123 – 131	7.5	86.2 – 93.1
Fort Myers Yacht Basin	2.0 – 8.6 [1.2 – 6.1]	3.6 – 6.6	106 – 126	5.0	83.3 – 91.5
Shell Point	17 – 34 [15 – 34]	4.5 – 7.6	25.5 – 124	1.4	83.2 – 90.8
McIntyre Creek	27.9 – 31.6 [30.3 – 33.9]	2.0 – 7.1	34.7 – 70.2	1.8 – 4.9	83.7 – 92.1
Tarpon Bay	30.4 – 34.5 [30.3 – 33.5]	3.3 – 8.5	13.2 – 33.3	0.9 – 2.4	83.7 – 91.1
Wulfert Flats	29.4 – 31.0 [30.1 – 34.2]	1.7 – 8.1	-----	5.6 – 100.1	84.9 – 92.3

Red values are outside of the preferred range.

<sup>a</sup> Salinity target values: BI < 5, FM < 10, SP = 10 – 30

<sup>b</sup> Dissolved O<sub>2</sub> target values: all sites > 4

<sup>c</sup> FDOM target values: BI < 70, FM < 70, SP < 11


<sup>d</sup> Chlorophyll target values: BI < 11, FM < 11, SP < 11

<sup>f</sup> Temperature target values: < 90


<sup>s</sup> Single sonde lower and surface layer or surface grab lab measurement

**Wildlife Impacts:** In the past week, the CROW wildlife hospital on Sanibel admitted 0 patients with suspected red tide/toxicosis.

**Shellfish Advisory:** Shellfish harvest area #6212 (Pine Island Sound Section 1) is **CLOSED** by the Florida Department of Agriculture and Consumer Services (FDACS) as of 6/10/24 as a precautionary closure due to increased HAB cell counts.



ACOE Daily Reports			
Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
6/04/24	2395	1659	2295
6/05/24	1831	1414	1956
6/06/24	1527	1300	2014
6/07/24	1514	1295	1889
6/08/24	1399	805	1201
6/09/24	1396	821	547
6/10/24	1765	1290	1126
7-day avg	1690	1226	1575



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.



Drift algae including *Hypnea*, *Agardhiella*, *Gracilaria* and *Halymenia* along the Western side of Sanibel Island on 6/7/24.



Water clarity at Lighthouse Beach Park on 6/10 at 1:40 PM on a rising tide (2.8 ft).