

MEMORANDUM

To: USACE Colonel Brandon L. Bowman, Major Cory Bell, Richard McMillen, SFWMD Governing Board,
Executive Director Drew Bartlett, Jennifer Reynolds, Lawrence Glenn, DEP Secretary Shawn Hamilton

From: Periodic Scientists Conference Call Participants

Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex

Holly Milbrandt & Dana Dettmar - City of Sanibel

Harry Phillips & Maya Robert - City of Cape Coral

Allie Pecenka, Rick Bartleson PhD & Matt Depaolis- Sanibel-Captiva Conservation Foundation

In coordination with Lee County

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **August 27- September 2, 2024**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of **2,263 cfs** at **S-79** with a 7-day average of **255 cfs (11%)** coming from the lake at **S-77**. **The 14-day moving average flow at S-79 is 2,213 cfs** and has been in the **stress flow envelope** (2,100- 2,600 cfs; RECOVER 2020) **for 8 days** after **23 days** in the **damaging flow envelope** (>2,600 cfs). **The 14-day moving average flow at S-77 was 213 cfs.**

Recommendation: With the onset of the rainy season and predictions for increased Atlantic storm intensity in the upcoming hurricane season, we ask the Army Corps to remain reactive to changing conditions in Lake Okeechobee and the Caloosahatchee River and estuary to support the ecological health of this system. In addition, we request the USACE manage flows to align with RECOVER 2020 optimum flow targets for the Caloosahatchee; being 750– 2,100 cfs as measured at S-79.

USACE Action: Lake Okeechobee stage is in the bottom third of Zone D (Zone D3 of the PA25 simulation) of the LOSOM regulation schedule, above the ecological envelope. The current climate outlook is for ENSO-neutral with La Niña favored to develop during September-November (ENSO- increased likelihood of below normal dry season rainfall north of the Lake). The District recommends USACE implements a non-harmful release from Lake Okeechobee to the Caloosahatchee Estuary with an average discharge of 2,000 cfs (7-day pulse) as measured at the S-79 structure, zero lake releases to the St. Lucie Estuary and zero lake releases to the Lake Worth Lagoon. The USACE should continue to track Red Tide and Blue Green Algae conditions, and should conditions change during this operational period, the USACE should look to reassess releases as needed.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was **4,837 AF** with **3,528 AF** to the Caloosahatchee through **S-77**, **-1 AF** to the St. Lucie canal through **S-308**, **1,310 AF** through the **L8 canal**, and **0 AF** to the EAA through **S-351**, **S-352**, and **S-354**. The total net inflow to the Lake was **28,634 AF** (**28,634 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1). Water conservation areas received flows of **2,969 AF**, **12,610 AF**, and **4,343 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **20,241 AF**.

*Data missing for S-310 from 8/27- 9/2, ENP from 8/27- 8/28 and Istokpoga, S-65E & S-65EX1 on 8/31.

Lake Level: 14.28 ft (Low Sub-Band)

Last Week: 14.16 ft

Last Year: 15.40 ft

7-Day Lake Recession Rate: +0.12 ft/week

Lake Okeechobee Inflow: 2,451 cfs

Lake Okeechobee Outflow: 87 cfs


Weekly Rainfall Total: WP Franklin: 1.26"

Ortona: 2.40"

Moore Haven: 4.12"

Cyanobacteria Status: On 9/3/24, sampling for cyanobacteria by the Lee County Environmental Lab reported no visible cyanobacteria across all sites.

Red Tide: On 8/30/24, the FWC reported that the red tide organism, *Karenia brevis*, was not observed in samples collected statewide over the past week.



Light Penetration				
Site	25% Iz	Target Values	Turbidity	Target Values
	meters		NTU	
Beautiful Is	0.6	> 1	2.0	< 18
Shell Point	1.6	>2.2	1.9	< 18
Causeway	2.9	> 2.2	1.5	< 5

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Upper Estuary Conditions: The 30-day average surface salinity at the Fort Myers Yacht Basin was 0.3 psu, within the suitable range for tape grass.

Lower Estuary Conditions: The average salinity at Shell Point RECON was 22 psu, in the optimal range for oysters but below optimal for seagrass.


Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	0.2 – 0.2 [0.2 – 0.2]	3.3 - 4.8	200	7.5	86.3– 91.5
Fort Myers Yacht Basin	0.3 – 1.2 [0.2 – 0.5]	ND	ND	ND	82.9– 87.4
Shell Point	8.4 - 30 [11 – 29]	ND	55	0.8	84.0– 89.0
McIntyre Creek	22.5 - 30 [27.1 – 23.5]	0.8 – 8.4	32.5 – 99	2.7 – 11.1	83.2 – 93.7
Tarpon Bay	21.6 - 32.1 [21.1 - 30.8]	2.1 – 8.9	23.1 – 73.3	1.1 – 8.1	83.4 – 90.5
Wulfert Flats	24.3 - 27.4 [25.3 - 27]	2.3 – 9.9	----	3.3 – 54.5	83.6 – 94.5


- Red values are outside of the preferred range.
- ^a Salinity target values: BI < 5, FM < 10, SP = 10 – 30
- ^b Dissolved O₂ target values: all sites > 4
- ^c FDOM target values: BI < 70, FM < 70, SP < 11
- ^d Chlorophyll target values: BI < 11, FM < 11, SP < 11
- ^f Temperature target values: < 90
- ^s Single sonde lower and surface layer or surface grab lab measurement

Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted 0 patients with suspected red tide/toxicosis.

Shellfish Advisory: Shellfish harvest area #6212 (Pine Island Sound Section 1; Aquaculture Leases) is **OPEN** while the public reef is **CLOSED** by the Florida Department of Agriculture and Consumer Services (FDACS) as of 7/13/24 due to presence of HAB *Pyrodinium bahamense*. SHA's 6222 (Pine Island Sound Sec. 2) and 6232 (Pine Island Sound Sec. 3) are **OPEN** as of 8/17/2024.



ACOE Daily Reports			
Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
8/27/24	2509	588	486
8/28/24	1769	751	490
8/29/24	1745	426	282
8/30/24	2496	738	34
8/31/24	2519	559	407
9/1/24	2077	643	87
9/2/24	2724	1155	0
7-day avg	2263	694	255



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 9/4/24 at 1:05 PM on a rising tide (2.8 ft).