

MEMORANDUM

To: USACE Colonel Brandon L. Bowman, Major Cory Bell, Richard McMillen, SFWMD Governing Board, Executive Director Drew Bartlett, Jennifer Reynolds, Lawrence Glenn, DEP Secretary Alexis Lambert

From: Periodic Scientists Conference Call Participants
Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex
Holly Milbrandt & Dana Dettmar - City of Sanibel
Allie Pecenka, Rick Bartleson PhD & Matt Depaolis- Sanibel-Captiva Conservation Foundation

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **March 4- 10, 2025**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of **1,974 cfs at S-79** with a 7-day average of **1,912 cfs (97%)** coming from the lake at **S-77**. **The 14-day moving average flow at S-79 was 2,097 cfs** and has been in the **optimum flow envelope** (750- 2,100 cfs; RECOVER 2020) for **1 day** after **18 days in the stress flow envelope** (2,100- 2,600 cfs; RECOVER 2020). **The 14-day moving average flow at S-77 was 1,677 cfs.**

Recommendation: We ask the USACE to structure recovery flows to the CRE in a format that will benefit the ecology of the ecosystems and align with RECOVER 2020 optimum flow targets of 750- 2,100 cfs measured at S-79.

USACE Action: Lake Okeechobee stage is in the middle third of Zone D (Zone D2 of the PA25 simulation) of the LOSOM regulation schedule. The current climate outlook is for La Niña and is expected to persist in the near-term. The District will continue to monitor conditions in the estuaries in anticipation of the onset of spawning season. The District recommends the USACE should continue non-harmful Recovery Operations for Lake Okeechobee as described in LOSOM while looking to implement potential reductions in flows as spawning is initiated this year based on estuarine conditions and climate forecasts. The USACE should continue to track Red Tide and Blue Green Algae conditions, and should conditions change during this operational period, the USACE should look to reassess releases as needed.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was **67,469 AF** with **23,120 AF** to the Caloosahatchee through **S-77**, **7,754 AF** to the St. Lucie canal through **S-308** and **36,595 AF** to the EAA through **S-351**, **S-352**, and **S-354**. The total net inflow to the Lake was **9,308 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1. Water conservation areas received flows of **55 AF**, **-898 AF**, and **7,126 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **1,855 AF**.

*Data missing from S-310 and L-8 from 3/4/25- 3/10/25, and missing from S-77 on 3/10/25.

Lake Level: 13.43 (Zone D2)

Last Week: 13.70 ft

Last Year: 15.94

7-Day Lake Recession Rate: -0.27 ft/week

Lake Okeechobee Inflow: 509 cfs

Lake Okeechobee Outflow: 4,920 cfs


Weekly Rainfall Total: WP Franklin: 0.03"

Ortona: 0.00"

Moore Haven: 0.11"

Cyanobacteria Status: On 3/10/25, sampling for cyanobacteria by the Lee County Environmental Lab reported **no visible cyanobacteria** across all sites.

Red Tide: On 2/28/25, the FWC reported that the red tide organism, *Karenia brevis*, was detected in 16 samples collected from Florida's Gulf Coast. In **Southwest Florida** over the past week, *K. brevis* was observed at background concentrations in Pinellas County, background to low concentrations in Hillsborough County, background to very low concentrations offshore of Manatee County, and background and very low concentrations in Charlotte County.



Light Penetration				
Site	25% Iz meters	Target Values	Turbidity NTU	Target Values
Beautiful Is	ND	> 1	ND	< 18
Shell Point	1.5	>2.2	1.7	< 18
Causeway	3.4	> 2.2	2.1	< 5

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Upper Estuary Conditions: The 30-day average surface salinity at the Fort Myers Yacht Basin was 2.8 psu, in the range for tape grass.

Lower Estuary Conditions: The weekly average salinity at the Shell Point RECON was 25 psu, in the optimal range for oysters but below optimal for seagrass. Phytoplankton biomass was low in SCCF's causeway and beach samples during the week.

Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	ND [0.2 - 0.5]	ND	ND	ND	ND
Fort Myers Yacht Basin	0.5- 4.0 [0.8- 5.1]	ND	ND	ND	67.7– 75.2
Shell Point	13 - 33 [16 – 32]	6.3 – 8.0	70	2.5	67.3– 74.6
McIntyre Creek	29.3 – 33.0 [27.4 – 30.7]	3.5 – 7.0	20.2—47.2	1.0 – 2.1	64.3 – 74.6
Tarpon Bay	28.0 – 34.6 [27.4 – 30.2]	5.7 – 9.4	13.7 – 71.6	0.8 – 3.2	66.2 – 74.0
Wulfert Flats	ND [ND]	ND	-----	ND	ND

Red values are outside of the preferred range.

^a Salinity target values: BI < 5, FM < 10, SP = 10 – 30

^b Dissolved O₂ target values: all sites > 4

^c FDOM target values: BI < 70, FM < 70, SP < 11


^d Chlorophyll target values: BI < 11, FM < 11, SP < 11

^f Temperature target values: < 90


^s Single sonde lower and surface layer or surface grab lab measurement
ND: no data

Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted **12 patients** with suspected red tide/toxicosis: 1 Adult Double-crested Cormorant (deceased), 1 Juvenile Double-crested Cormorant (still in care), 1 Adult White Ibis (still in care), 1 Adult Laughing Gull (deceased), and 8 Royal Terns: 1 Juvenile (deceased), 3 adults (deceased) and 4 adults (still in care).

Shellfish Advisory: Shellfish harvest area #6212 (Pine Island Sound Section 1; Aquaculture Lease and Public Reef) is **CLOSED** due to the presence of *Karenia brevis* as of 11/06/24. SHA #6222 (North Matlacha Pass) and SHA #6232 (South Matlacha Pass) are **CLOSED** by the Florida Department of Agriculture and Consumer Services (FDACS) due to the presence of *Karenia brevis* as of 1/30/25.



ACOE Daily Reports			
Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
3/4/2025	2196	1920	2124
3/5/2025	1926	1829	1938
3/6/2025	1993	1552	1872
3/7/2025	1697	1429	1964
3/8/2025	1744	1360	1884
3/9/2025	2066	1365	1864
3/10/2025	2196	1668	1737
7-day avg	1974	1589	1912



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 3-12-25 at 2:00 PM on a falling tide (1.7 ft).