

MEMORANDUM

To: USACE Colonel Brandon L. Bowman, Major Cory Bell, Richard McMillen, SFWMD Governing Board,
Executive Director Drew Bartlett, Jennifer Reynolds, DEP Secretary Alexis Lambert

From: Periodic Scientists Conference Call Participants

Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex

Holly Milbrandt & Dana Dettmar - City of Sanibel

Allie Pecenka, Rick Bartleson PhD & Matt Depaolis- Sanibel-Captiva Conservation Foundation

With contributions from Harry Phillips & Maya Robert PhD- City of Cape Coral

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **May 6- 12, 2025**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of **348 cfs** at **S-79** with a 7-day average of **496 cfs** (100%) coming from the lake at **S-77**. **The 14-day moving average flow at S-79 was 345 cfs** and has been **below the optimum flow envelope** (<750 cfs) for **15 days** after being in the optimum flow envelope (750- 2,100 cfs; RECOVER 2020) for **47 days**. **The 14-day moving average flow at S-77 was 691 cfs.**

Recommendation: We ask that S-79 be managed to ensure actual and projected flows are aligning, as actual flows have been consistently lower than the constant 500 cfs flow schedule, driving weekly averages under the MFL and creating damaging salinity levels. The current flow target at S-79 is below the RECOVER 2020 optimum flow envelope and salinities are continuing to rise quickly in the Caloosahatchee estuary, threatening the health of indicator species like oysters and tape grass. Damages to these populations take decades to recover from, including oysters which are currently in a critical spawning period. The CRE is in a vulnerable position and cannot withstand flows being dropped any lower. We ask that flows at S-79 be within the optimum envelope of 750-2,100 cfs to preserve the health of the estuary and all species inhabiting it.

USACE Action: Lake Okeechobee stage is in the lower portion of Zone D (Zone D3 of the PA25 simulation) of the LOSOM regulation schedule. ENSO-neutral conditions are present and is favored during the summer. The District continues to monitor conditions in the estuaries, as well as the systemwide conditions, as the dry season progresses. As such, the District recommends that USACE should continue non-harmful Recovery Operations for Lake Okeechobee as described in LOSOM recognizing that there is a higher chance to achieve Recovery targets. To maintain favorable salinity levels in the estuaries and begin to conserve water, it is recommended that flow targets for the Caloosahatchee Estuary should ensure the delivery of the Minimum Flow and Level via a targeted steady release of 500 cfs and operational changes at S-77, S-78, and S-79 should be implemented to ensure the 500 cfs is delivered on a daily basis; flow targets for the St. Lucie Estuary should remain at 0 cfs, and the flow target for the Lake Worth Lagoon should remain at 0 cfs. The District will continue to monitor salinity conditions in the estuaries and water supply conditions throughout the system as the dry season progresses to assess future operational recommendations.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was **24,339 AF** with **6,176 AF** to the Caloosahatchee through **S-77**, **1,214 AF** to the St. Lucie canal through **S-308** and **16,949 AF** to the EAA through **S-351**, **S-352**, and **S-354**. The total net inflow to the Lake was **1,613 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1. Water conservation areas received flows of **1,201 AF**, **2,392 AF**, and **2,230 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **147 AF**.

*Data missing from S-310 and L-8 on 5/6/25- 5/12/25, and from S-77, S-78, S-79, S-351, S-352, S-354, S-308 and S-80 on 5/12/25.

Lake Level: 10.98 ft (Zone D3)

Last Week: 11.17 ft

Last Year: 13.77 ft

7-Day Lake Recession Rate: -0.19 ft/week

Lake Okeechobee Inflow: 115 cfs


Lake Okeechobee Outflow: 474 cfs

Weekly Rainfall Total: WP Franklin: 2.37"

Ortona: 3.59"

Julian Keen Jr.: 4.23"

Cyanobacteria Status: On 5/5/25, sampling for cyanobacteria by the Lee County Environmental Lab reported **abundant** streaks and some accumulation of primarily *Dolichospermum* and some *Microcystis* at the **Alva Boat Ramp**. Some specks of cyanobacteria were observed **upstream of the Franklin Locks**.



Light Penetration

Site	25% Iz meters	Target Values	Turbidity NTU	Target Values
Fort Myers	1.2	> 1	4.5	< 18
Shell Point	2.1	> 2.2	2.2	< 18
Causeway	5.3	> 2.2	1.3	< 5

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Red Tide: On 5/9/25, the FWC reported that the red tide organism, *Karenia brevis*, was detected at background concentrations in two samples from Florida's Gulf Coast; in Sarasota County and Bay County.

Upper Estuary Conditions: The 30-day average surface salinity at the Fort Myers Yacht Basin was 6.7 psu, in the range for tape grass and the weekly average rose to 12 psu.

Lower Estuary Conditions: The weekly average salinity at the Shell Point RECON was **33 psu**, in the optimal range for seagrass, but **above optimal for oysters**. *Pseudo-nitzschia* concentrations in samples from the causeway and Pine Island Sound were < 25,000 cells/L and small diatoms were dominant.

Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	4.2 – 6.0 [3.3 – 6.1]	3.8 – 6.6	115	7.5	82.4 – 88.3
Fort Myers Yacht Basin	9.3 - 15 [6.1 – 12]	4.2– 6.7	73	3.0	79.6 – 88.3
Shell Point	27 – 36 [25 – 36]	4.6 – 7.8	25	1.7	80.9 – 86.4
McIntyre Creek	33.3 – 36.3 [33.9 – 35.2]	1.5 – 5.9	12.3 – 36.5	0.8 – 2.9	80.6 – 89.1
Tarpon Bay	33.9 – 36.7 [34.7 – 36.2]	3.9 – 8.8	8.5 – 20.6	0.7 – 2.2	81.2 – 87.3
Wulfert Flats	34.3 – 36.2 [34.5 – 35.4]	2.9 – 9.5	-----	3.1 – 14.1	81.5 – 88.5

Red values are outside of the preferred range.

^a Salinity target values: BI < 5, FM < 10, SP = 10 – 30

^b Dissolved O₂ target values: all sites > 4

^c FDOM target values: BI < 70, FM < 70, SP < 11

^d Chlorophyll target values: BI < 11, FM < 11, SP < 11


^f Temperature target values: < 90

^s Single sonde lower and surface layer or surface grab lab measurement

ND: no data


Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted **0 patients** with suspected red tide/toxicosis.

Shellfish Advisory: Shellfish harvest area #6212 (Pine Island Sound Section 1; Aquaculture Lease and Public Reef) is **OPEN** by the Florida Department of Agriculture and Consumer Services (FDACS) as of 4/3/25. SHA #6222 (North Matlacha Pass) is **OPEN** as of 4/15/25. SHA #6232 (South Matlacha Pass) is **OPEN** as of 3/21/25.



ACOE Daily Reports

Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
5/6/25	277	344	704
5/7/25	356	343	572
5/8/25	396	341	572
5/9/25	382	205	568
5/10/25	328	148	380
5/11/25	314	250	179
5/12/25	383	296	NR
7-day avg	348	275	496



Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 5-13-25 at 2:15 PM on a falling tide (2.9 ft).