

MEMORANDUM

To: USACE Colonel Brandon L. Bowman, Major Cory Bell, Richard McMillen, SFWMD Governing Board,
Executive Director Drew Bartlett, Jennifer Reynolds, DEP Secretary Alexis Lambert

From: Periodic Scientists Conference Call Participants
Kevin Godsea & Avery Renshaw - J.N. "Ding" Darling National Wildlife Refuge (NWR) Complex
Holly Milbrandt & Dana Dettmar - City of Sanibel
Allie Pecenka, Rick Bartleson PhD & Matt DePaolis- Sanibel-Captiva Conservation Foundation
With contributions from Harry Phillips & Maya Robert PhD- City of Cape Coral

Subject: Caloosahatchee & Estuary Conditions Report

Reporting Period: **January 27- February 2, 2026**

This report provides a scientific assessment of Caloosahatchee River and Estuary conditions and how these conditions affect the health, productivity, and function of the system.

Caloosahatchee Conditions Summary: Flow to the Caloosahatchee Estuary had a 7-day average of **419 cfs** at **S-79** with a 7-day average of **755 cfs** coming from the lake at **S-77**. **The 14-day moving average flow at S-79 was 401 cfs** and has been **below the optimum flow envelope (<750 cfs) for 100 days**. **The 30-day moving average flow at S-79 has been below 457 cfs for 67 consecutive days**. **The 14-day moving average flow at S-77 was 641 cfs**.

Recommendation: The 14-day moving average flow at S-79 has remained below the optimum flow envelope (750- 2,100 cfs) for the past **100 days** and the 30-day moving average at S-79 has been **below 457 cfs for 67 consecutive days**. Salinity levels have risen above the optimum range for tape grass (<10 psu) in the upper estuary and above the optimum range for oysters (10-25 psu) in the lower estuary (RECOVER 2020). We recommend that supplemental releases from S-77 be made to ensure S-79 flows remain above a targeted steady release of at least 750 cfs to maintain necessary salinity levels in the Caloosahatchee estuary.

USACE Action: On January 25 the daily average Lake Okeechobee stage was 11.68 feet NAVD88 (13.09 feet NGVD29), which placed it within the lower portion of Zone D (Zone D3 of the PA25 simulation) of the Lake Okeechobee System Operating Manual (LOSOM). Lake stage decreased by 0.10 feet over the preceding 7-day period. There is a 75% chance of a transition to ENSO-neutral during January-March 2026. The District will continue to monitor conditions in the estuaries, as well as the systemwide conditions. With the initiation of the dry season, Normal Lake Operations continue pursuant to the considerations in LOSOM as informed by PA25. It is recommended that flow targets for the Caloosahatchee Estuary should rely on basin flows to ensure the delivery of the Minimum Flow and Level, but use Lake Okeechobee flows from S-77 to ensure S-79 flows remain above a targeted steady release of 350 cfs; flow targets for the St. Lucie Estuary and Lake Worth Lagoon should remain at 0 cfs consistent with Normal Operations within Zone D.

Lake Flows: In the past 7 days the total outflow from Lake Okeechobee was **56,513 AF** with **10,542 AF** to the Caloosahatchee through **S-77**, **1,574 AF** to the St. Lucie canal through **S-308** and **44,397 AF** to the EAA through **S-351**, **S-352**, and **S-354**. The total net inflow to the Lake was **4,093 AF** from Fisheating Creek, S-71, S-72, S-84s, S-65EX, and S-65EX1. Water conservation areas received flows of **143 AF**, **85 AF**, and **1,471 AF** at **WCA1**, **WCA2**, and **WCA3**, respectively. Everglades National Park received **234 AF**.

*Data missing from S-310 and L-8 from 1/27- 2/2, from S-79 on 2/2 and 1/26, and from S-80 on 1/28 and 2/2.

Lake Level: 12.72 (Zone D3)

Last Week: 12.98 ft

Last Year: 14.36 ft

7-Day Lake Recession Rate: -0.26 ft/ wk.

Lake Okeechobee Inflow: 256 cfs

Lake Okeechobee Outflow: 4,470 cfs


Weekly Rainfall Total: WP Franklin: 0.10"

Ortona: 0.02"

Julian Keen Jr.: 0.01"

Red Tide: On 1/30/26, the FWC reported that the red tide organism, *Karenia brevis*, was [observed in 12 samples from Florida](#). In Southwest Florida over the past week, *K. brevis* was observed at background concentrations in one sample collected offshore of Hillsborough County, and background and very low concentrations in two samples collected offshore of Lee County.

Cyanobacteria Status: On 2/2/26, sampling for cyanobacteria by the Lee County Environmental Lab reported *Microcystis*, *Dolichospermum* and cyano filaments as **moderately abundant** at the **Alva Boat ramp** appearing as **streaks** and **abundant upstream of the Franklin Locks** and at the **Davis Boat Ramp**, appearing as **streaks with accumulation**.



Light Penetration				
Site	25% Iz	Target Values	Turbidity	Target Values
	meters		NTU	
Fort Myers	1.0	> 1	2.5	< 18
Shell Point	1.7	>2.2	3.0	< 18
Causeway	2.6	> 2.2	9.0	< 5

25% Iz is the depth (z) where irradiance (I) is 25% of surface irradiance. Target values indicate the depth of light penetration needed for healthy seagrass.

Upper Estuary Conditions: The 30-day moving average surface salinity at the Fort Myers Yacht Basin has been over 10 psu for **67 days** and was **15 psu**, above the suitable range for **tape grass**. The weekly average was 14 psu.

Lower Estuary Conditions: The weekly average salinity at the Shell Point RECON was **32 psu**, in the optimal range for seagrass but **above optimal for oysters**. The phytoplankton in SCCF samples from Sanibel beaches during the week included mainly diatoms, low densities of *Pseudo-nitzschia* and no *Karenia* cells.

Water Quality Conditions:

Monitor Site	Salinity (psu) ^a [previous week]	Diss O ₂ (mg/L) ^b	FDOM (qsde) ^c	Chlorophyll (µg/L) ^d	Temperature (°F)
Beautiful Island	7.3 – 9.9 [6.9 – 9.7]	6.4 – 9.5	145	6.3	59.3 - 78.4
Fort Myers Yacht Basin	11 – 19 [11 – 20]	7.1 – 9.1	100	4.5	52.1– 72.8
Shell Point	24 – 34 [25 – 34]	6.4 - 8.6	40	2.3	53.1 – 70.5
McIntyre Creek	34.4 – 36.1 [34.0 – 35.2]	5.0 – 9.7	11.5 – 31.0	1.1 – 2.4	47.6 – 67.3
Tarpon Bay	33.3 – 35.9 [33.7 – 35.3]	6.8 – 8.9	15.4 – 51.0	0.8 – 4.0	50.0 – 66.2
Wulfert Flats	---- [----]	----	----	----	----

Red values are outside of the preferred range.

^a Salinity target values: BI < 5, FM < 10, SP = 10 – 30

^b Dissolved O₂ target values: all sites > 4

^c FDOM target values: BI < 70, FM < 70, SP < 11


^d Chlorophyll target values: BI < 11, FM < 11, SP < 11

^f Temperature target values: < 90


^s Single sonde lower and surface layer or surface grab lab measurement
ND: no data

Wildlife Impacts: In the past week, the CROW wildlife hospital on Sanibel admitted 1 patient for suspected red tide/toxicosis: 1 adult cattle egret (deceased).

Shellfish Advisory: Shellfish harvest area #6212 (Pine Island Sound Section 1); Aquaculture Lease and Public Reef is **OPEN** by the Florida Department of Agriculture and Consumer Services (FDACS) as of 9/24/25. SHA #6222 (North Matlacha Pass) and SHA #6232 (South Matlacha Pass) are **OPEN** as of 12/6/25.



USACE Daily Reports			
Date	S79 Flow (cfs)	S78 Flow (cfs)	S77 Flow (cfs)
1/27/26	568	214	806
1/28/26	473	174	576
1/29/26	336	250	623
1/30/26	305	318	878
1/31/26	465	237	800
2/1/26	425	147	855
2/2/26	358	152	750
7-day avg	419	213	755




Daily average bottom salinity data for the last 14-days from sampling locations within the tidal Caloosahatchee River Estuary relative to oyster health (Sanibel, Shell Point and Cape Coral) and tape grass (*Vallisneria americana*) health (Ft. Myers only) conditions.

*Ft. Myers sensor is in the lower strata



Water clarity at Lighthouse Beach Park on 2-3-26 at 1:55 PM on a slack tide (1.5 ft).



Cyanobacteria accumulation at the Davis Boat Ramp on 2/3/26 at 2:15 pm.



Cold-stunned and deceased marine species in the wrack line on Sanibel Island beaches over the past week.

